

CHEM**ELECTRO**CHEM

Supporting Information

© Copyright Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, 2014



Proof of Concept of the Electrochemical Sensing of 3-Iodothyronamine (T₁AM) and Thyronamine (T₀AM)

Luís M. Gonçalves,^{*,[a, b, c]} Manuela M. Moreira,^[a, d] Carla F. Azevedo,^[a] Inês M. Valente,^[a]
João C. Sousa,^[b, c] Thomas S. Scanlan,^[e] Richard G. Compton,^[f] and José A. Rodrigues^[a]



celc_201402165_sm_miscellaneous_information.pdf

Supporting information

I – a) current vs scan rate and b) current vs square rate of scan rate




0.153 mM of T_0AM on a GCE




0.174 mM of T_1AM on a GCE

II – Benzoquinone



Blue is the 10th scan of the figure below and red is benzoquinone. Both are done at 1000 mV/s.



Cyclic voltammograms of aqueous phosphate buffer solution (pH 7) containing 0.174 mM of T₁AM on a GCE, cyclic voltammetry was run between 0.0 and +1.0 V vs. Ag|AgCl at different scan rates. Sequential cyclic voltammograms at 1000 mV/s showing the appearance of a reductive and an oxidative peak around 0.0 and +0.2 V vs. Ag|AgCl, respectively.